

TREATMENT OF RHEUMATOID ARTHRITIS WITH CHIMERIC MONOCLONAL ANTIBODIES TO TUMOR NECROSIS FACTOR α

MICHAEL J. ELLIOTT, RAVINDER N. MAINI, MARC FELDMANN, ALICE LONG-FOX, PETER CHARLES, PETER KATSIKIS, FIONULA M. BRENNAN, JEAN WALKER, HANNY BIJL, JOHN GHRAYEB, and JAMES N. WOODY

Objective. To evaluate the safety and efficacy of a chimeric monoclonal antibody to tumor necrosis factor α (TNF α) in the treatment of patients with rheumatoid arthritis (RA).

Methods. Twenty patients with active RA were treated with 20 mg/kg of anti-TNF α in an open phase I/II trial lasting 8 weeks.

Results. The treatment was well tolerated, with no serious adverse events. Significant improvements were seen in the Ritchie Articular Index, which fell from a median of 28 at study entry to a median of 6 by week 6 ($P < 0.001$), the swollen joint count, which fell from 18

to 5 ($P < 0.001$) over the same period, and in the other major clinical assessments. Serum C-reactive protein levels fell from a median of 39.5 mg/liter at study entry to 8 mg/liter at week 6 ($P < 0.001$), and significant decreases were also seen in serum amyloid A and interleukin-6 levels.

Conclusion. Treatment with anti-TNF α was safe and well tolerated and resulted in significant clinical and laboratory improvements. These preliminary results support the hypothesis that TNF α is an important regulator in RA, and suggest that it may be a useful new therapeutic target in this disease.

From the Clinical Immunology and Sunley Divisions, Kennedy Institute of Rheumatology, and the Academic Department of Rheumatology, Charing Cross and Westminster Medical School, London, United Kingdom; and Centocor Inc., Malvern, Pennsylvania.

Supported in part by grants from the Arthritis and Rheumatism Council of Great Britain. Dr. Elliott's work was supported by the Michael Mason Fellowship from the Arthritis Foundation of Australia, and by the DEV Starr Fellowship from the Royal Australasian College of Physicians.

Michael J. Elliott, MB, BS, PhD, FRACP: Clinical Immunology Division, Kennedy Institute of Rheumatology; Ravinder N. Maini, MB, BChir, FRCP: Clinical Immunology Division, Kennedy Institute of Rheumatology; Marc Feldmann, MB, BS, PhD, FRCP: Sunley Division, Kennedy Institute of Rheumatology; Alice Long-Fox, RGN: Clinical Immunology Division, Kennedy Institute of Rheumatology; Peter Charles, FIMLS: Department of Rheumatology, Charing Cross and Westminster Medical School; Peter Katsikis, MD, PhD: Sunley Division, Kennedy Institute of Rheumatology; Fionula M. Brennan, PhD: Sunley Division, Kennedy Institute of Rheumatology; Jean Walker, RGN, RM: Clinical Immunology Division, Kennedy Institute of Rheumatology; Hanny Bijl, MD: Centocor, Inc; John Ghrayeb, PhD: Centocor Inc; James N. Woody, MD, PhD: Centocor, Inc.

Address reprint requests to Michael J. Elliott, PhD, Kennedy Institute of Rheumatology, 6 Bute Gardens, London W6 7DW, United Kingdom.

Submitted for publication May 28, 1993; accepted in revised form August 10, 1993.

Despite optimal use of current antirheumatic therapy, the outcome for many patients with rheumatoid arthritis (RA) consists of pain, disability, and premature death (1-3). As a response to the need for more effective and less toxic treatment, and to an increase in our understanding of the pathogenic mechanisms in RA, several groups have used monoclonal antibodies as therapeutic agents in this disease (4-10). Such immunotherapy has been, in most cases, targeted specifically to the T cell, a strategy based on evidence that T cells are involved in the initiation and maintenance of RA (11).

Here, we outline an alternative immunotherapeutic strategy, which involves the use of monoclonal antibodies with specificity for a cytokine, tumor necrosis factor α (TNF α). This approach is based on a body of knowledge regarding the role of cytokines in general, and of TNF α in particular, in the inflammatory process in RA. The first clearly documented study demonstrated the presence of interleukin-1 (IL-1) in RA synovial fluid (12). Subsequently, we and others have reported the presence and local synthesis in

BEST AVAILABLE COPY

EXHIBIT

F

Table 1. Demographic features of 20 patients with refractory rheumatoid arthritis

Patient	Age/sex	Disease duration (years)	Previous DMARDs*	Concomitant therapy†
1	48/F	7	SSZ, DP, GST, AUR, MTX, AZA, HCQ	Pred. 5 mg
2	63/F	7	SSZ, GST, DP	Para. 1-2 gm
3	59/M	3	AUR, HCQ, GST, MTX, SSZ	Pred. 10 mg, Indo. 225 mg
4	56/M	10	GST, DP, AZA, SSZ	Pred. 12.5 mg, Ibu. 2 gm, Para. 1-2 gm
5	28/F	3	GST, SSZ, DP, AZA	Pred. 8 mg, Para. 1-2 gm, Code. 16 mg
6	40/M	3	SSZ, HCQ, AUR	Nap. 1 gm
7	54/F	7	DP, GST, SSZ, AZA, MTX	Para. 1-2 gm, Code. 16-32 mg
8	23/F	11	HCQ, GST, SSZ, MTX, AZA	Pred. 7.5 mg, Dicl. 100 mg, Para. 1-2 gm, Dext. 100-200 mg
9	51/F	15	GST, HCQ, DP, MTX	Pred. 7.5 mg, Dicl. 125 mg, Para. 1-3 gm
10	47/F	12	SSZ, CYC, MTX	Ben. 4 gm
11	34/F	10	DP, SSZ, MTX	Pred. 10 mg, Para. 1.5 gm, Code. 30-90 mg
12	57/F	12	GST, MTX, DP, AUR	Asp. 1.2 gm
13	51/F	7	SSZ, AZA	Para. 1-4 gm
14	72/M	11	GST, DP, AZA, MTX	Pred. 5 mg, Para. 1-4 gm, Code. 16-64 mg
15	51/F	17	HCQ, DP, SSZ, MTX	Asp. 0.3 gm
16	62/F	16	GST, DP, SSZ, MTX, AZA	Para. 1-4 gm, Code. 16-64 mg
17	56/F	11	SSZ, DP, GST, MTX, HCQ, AZA	Pred. 7.5 mg, Eto. 600 mg, Para. 1-2 gm, Dext. 100-200 mg
18	48/F	14	GST, MTX, DP, SSZ, AUR, AZA	Pred. 7.5 mg, Indo. 100 mg, Para. 1-3 gm
19	42/F	3	SSZ, MTX	Fen. 450 mg, Ben. 6 gm, Code. 30 mg
20	47/M	20	GST, DP, SSZ, AZA	Pred. 10 mg, Para. 1-3 gm

* Disease-modifying antirheumatic drugs (DMARDs) were SSZ = sulfasalazine; DP = D-penicillamine; GST = gold sodium thiomaleate; AUR = auranofin; MTX = methotrexate; AZA = azathioprine; HCQ = hydroxychloroquine; CYC = cyclophosphamide.

† Daily doses are shown. Pred. = prednisolone; Para. = paracetamol; Indo. = indomethacin; Ibu. = ibuprofen; Code. = codeine phosphate; Nap. = naprosyn; Dicl. = diclofenac; Dext. = dextropropoxyphene; Ben. = benorylate; Asp. = aspirin; Eto. = etodolac; Fen. = fenbufen.

rheumatoid synovial membrane of many cytokines, including IL-1 (13), TNF α (13,14), IL-6 (15), granulocyte-macrophage colony-stimulating factor (GM-CSF; 16), IL-8 (17), and transforming growth factor β (TGF β) (18,19).

We have investigated the relationships between these cytokines in RA, using a synovial culture system in which dissociated rheumatoid synovial cells are allowed to spontaneously re-aggregate in vivo. Even in the absence of extrinsic stimulation, such cells express high levels of cytokines and HLA class II molecules (20). Using this system, we showed that production of bioactive IL-1 was abrogated by neutralizing antibodies to TNF α , but not by antibodies to TNF β or by normal rabbit IgG (21). This occurred in rheumatoid, but not osteoarthritic, cultures and suggested to us that TNF α was of particular importance as a regulatory cytokine. Subsequent analysis reinforced this concept, with the demonstration that another proinflammatory cytokine, GM-CSF, was regulated in the synovial membrane by TNF α (22) and that TNF α receptor expression, necessary for transmitting TNF α signals, was up-regulated in rheumatoid synovium (23,24).

Two recent mouse studies provide further insight into the importance of TNF α in arthritis. Keffer et al (25) described a mouse transgenic for the human TNF α gene, which expressed high levels of human TNF α in vivo and which reproducibly developed arthritis beginning at 4 weeks of age. The disease in these animals could be prevented by administration of monoclonal antibodies to human TNF α . In separate experiments in our own laboratory, we showed that in the type II collagen arthritis model in the DBA/1 mouse, the hamster anti-murine TNF monoclonal antibody TN3.19.2 significantly ameliorated the inflammation and tissue destruction when administered before or after the onset of disease (26).

Based on these considerations, it was of interest to determine the effect of therapy with a chimeric (human IgG1, murine Fv) monoclonal antibody to human TNF α in patients with rheumatoid arthritis. We report here that anti-TNF α therapy was safe and well tolerated, and induced marked improvements in both clinical and laboratory disease measures. These findings are consistent with our postulate concerning the critical role of TNF α in the pathogenesis of RA (27,28), and suggest that TNF α may be a useful therapeutic target in this disease.

Table 3. Changes in laboratory measures following treatment of rheumatoid arthritis patients with cA2*

Week of trial	Hgb, gm/liter	WBC, $\times 10^9$ /liter	Platelets, $\times 10^9$ /liter	ESR, mm/hour	CRP, mg/liter	SAA, mg/ml	RF, inverse liter
Screen	117, 98-146	7.9, 3.9-15.2	352, 274-631	59, 18-87	42, 9-107	ND	ND
0	113, 97-144	9.0, 4.9-15.7	341, 228-710	55, 15-94	39.5, 5-107	245, 18-1,900	2,560, 160-10,240
1	114, 96-145 (>0.05)	8.5, 3.6-13.6 (>0.05)	351, 223-589 (>0.05)	26, 13-100 (>0.05)	5, 0-50 (<0.001 †)	58, 0-330 (<0.001 ; <0.003 †)	ND
2	112, 95-144 (>0.05)	8.2, 4.3-12.7 (>0.05)	296, 158-535 (<0.04 ; >0.05 †)	27, 10-90 (<0.02 ; >0.05 †)	5.5, 0-80 (<0.001 ; <0.003 †)	80, 11-900 (<0.02 ; <0.04 †)	ND
3	110, 89-151 (>0.05)	9.0, 3.7-14.4 (>0.05)	289, 190-546 (<0.03 ; >0.05 †)	27, 12-86 (<0.04 ; >0.05 †)	7, 0-78 (<0.001 ; <0.002 †)	ND	ND
4	112, 91-148 (>0.05)	8.2, 4.7-13.9 (>0.05)	314, 186-565 (>0.05)	23, 10-87 (<0.04 ; >0.05 †)	10, 0-91 (<0.004 ; <0.02 †)	ND	ND
6	116, 91-159 (>0.05)	9.1, 2.9-13.9 (>0.05)	339, 207-589 (>0.05)	23, 12-78 (<0.03 ; >0.05 †)	8, 0-59 (<0.001 †)	ND	ND
8	114, 94-153 (>0.05)	7.6, 4.2-13.5 (>0.05)	339, 210-591 (>0.05)	30, 7-73 (>0.05)	6, 0-65 (<0.001 †)	ND	480, 40-5,120 (>0.05)

* Values are the median, range (P) for 20 patients for the initial screen and weeks 0-2, and for 19 patients thereafter. Patient 15 dropped out after week 2 of study. For rheumatoid factor (RF), only those patients with week 0 titers $\geq 1:160$ in the particle agglutination assay were included ($n = 14$). All P values versus week 0, by Mann-Whitney test. Normal ranges: hemoglobin (Hgb) 120-160 gm/liter in females and 135-175 gm/liter in males; white blood cell (WBC) count $4-11 \times 10^9$ /liter; platelet count $150-400 \times 10^9$ /liter; erythrocyte sedimentation rate (ESR) <15 mm/hour in females and <10 mm/hour in males; C-reactive protein (CRP) <10 mg/liter; serum amyloid A (SAA) <10 mg/ml. ND = not done.

questioned concerning possible adverse events before each infusion and at weeks 1, 2, 3, 4, 6, and 8 of the trial. A complete physical examination was performed at screening and at week 8. In addition, patients were monitored by standard laboratory tests including a complete blood cell count, and levels of C3 and C4 components of complement, IgG, IgM, and IgA; serum electrolytes, creatinine, urea, alkaline phosphatase, aspartate transaminase, and total bilirubin.

Sample times for these tests were between 0800 and 0900 hours (preinfusion) and 1200-1400 hours (24 hours postinfusion). Blood tests subsequent to day 1 were performed in the morning, usually between 0700 and 1200 hours. Urine analysis and culture were also performed at each assessment point.

Response assessment. The patients were assessed for response to cA2 at weeks 1, 2, 3, 4, 6, and 8 of the trial. The assessments were all made between 0700 and 1300 hours by the same observer (AL-F). The following clinical assessments were made: duration of morning stiffness (minutes), pain score (0-10 cm on a visual analog scale), Ritchie Articular Index (maximum score 69) (32), number of swollen joints (28 joint count) (validation described in ref. 33), grip strength (0-300 mm Hg, mean of 3 measurements per hand, by sphygmomanometer cuff), and an assessment of function (the Stanford Health Assessment Questionnaire [HAQ], modified for British patients [34]). In addition, the patients' global assessments of response were recorded using a 5-point scale (worse, no response, fair response, good response, excellent response).

Routine laboratory indicators of disease activity included complete blood cell counts, C-reactive protein (CRP) levels (by rate nephelometry), and the erythrocyte sedimentation rate (ESR; Westergren). Followup assessments were made at monthly intervals after the conclusion of the formal trial period, in order to assess the duration of response.

Analysis of improvement in individual patients was made using two separate indices. First, an IDA was calculated for each time point according to the method of Mallya and Mace (30), with input variables of morning stiffness, pain score, Ritchie Articular Index, grip strength, ESR, and Hgb. The second index calculated was that of Paulus et al (35), which uses input variables of morning stiffness, ESR, joint pain/tenderness, joint swelling, and patient's and physician's global assessments of disease severity.

To calculate the presence (or otherwise) of a response according to this index, two approximations were made to accommodate our data. The swollen joint count used by us (nongraded total of swollen joints of 28 joints assessed), which has been validated (33), was used in place of the more extensive graded count described by Paulus et al, and the patient's and physician's global assessments of response recorded by us were approximated to the global assessments of disease activity used by Paulus et al (35). In addition to determining response according to these published indices, we selected 6 disease activity assessments of interest (morning stiffness, pain score, Ritchie Articular Index, swollen joint count, ESR, and CRP) and calculated their mean percentage improvement. We have used this value to give an indication of the degree of improvement seen in responding patients.

Immunologic investigations. Rheumatoid factors were measured using the rheumatoid arthritis particle agglutination assay (RAPA) (FujiBerio Inc, Tokyo, Japan), in which titers of 1:160 or greater were considered significant. Rheumatoid factor isotypes were measured by enzyme-linked immunosorbent assay (ELISA) (Cambridge Life Sciences, Ely, UK). Addition of cA2, at concentrations of up to 200 μ g/ml, to these assay systems did not alter the assay results (data not shown).

Antinuclear antibodies were detected by immunoflu-

orescence on HEp-2 cells (Biodiagnostics, Upton, UK), and antibodies to extractable nuclear antigens were measured by counterimmunoelectrophoresis with polyantigen extract (Biodiagnostics). Sera positive by immunofluorescence were also screened for antibodies to DNA by the Farr assay (Kodak Diagnostics, Amersham, UK). Anticardiolipin antibodies were measured by ELISA (Shield Diagnostics, Dundee, Scotland). Serum amyloid A (SAA) was measured by sandwich ELISA (Biosource International, Camarillo, CA). Antiglobulin responses to the infused chimeric antibody were measured by an in-house ELISA, using cA2 as a capture reagent.

Cytokine assays. Bioactive TNF was measured in sera using the WEHI 164 clone 13 cytotoxicity assay (36). Total IL-6 was measured in sera using a commercial immunoassay (Medgenix Diagnostics, Brussels, Belgium) and using a sandwich ELISA developed in-house, with monoclonal antibodies provided by Dr. F. di Padova (Basel, Switzerland). Microtiter plates were coated with monoclonal antibody LNI 314-14 at a concentration of 3 μ g/ml for 18 hours at 4°C, and blocked with 3% bovine serum albumin in 0.1M phosphate buffered saline, pH 7.2. Undiluted sera or standards (recombinant HuIL-6, 0–8.1 μ g/ml) were added to the wells in duplicate and incubated for 18 hours at 4°C. Bound IL-6 was detected by incubation with monoclonal antibody LNI 110-14 for 90 minutes at 37°C, followed by biotin-labeled goat anti-murine IgG2b for 90 minutes at 37°C (Southern Biotechnology, Birmingham, AL). The assay was developed using streptavidin-alkaline phosphatase (Southern Biotechnology) and *p*-nitrophenyl phosphate as a substrate, and the optical density read at 405 nm.

Statistical analysis. Data for week 0 versus subsequent time points were compared for each assessment using the Mann-Whitney test. For comparison of rheumatoid factor titers (by RAPA), the data were expressed as dilutions before applying the test.

This was an exploratory study, in which prejudices about the optimal times for assessment were not possible. Although it has not been common practice to adjust for multiple statistical comparisons in such studies (4–10), a conservative statistical approach would require adjustment of *P* values to take into account analysis at several time points. The *P* values have therefore been presented in two forms: unadjusted, and after making allowance for analysis at multiple time points by use of the Bonferroni adjustment. Where *P* values remained <0.001 after adjustment, a single value only is given. A *P* value of <0.05 is considered significant.

RESULTS

Safety of cA2. The administration of cA2 was exceptionally well tolerated, with no headache, fever, hemodynamic disturbance, allergy, or other acute manifestation. No serious adverse events were recorded during the 8-week trial. Two minor infective episodes were recorded, each "possibly related" to cA2: patient 15 presented at week 2 with clinical features of bronchitis. Sputum culture grew only nor-

mal commensals. She had a history of smoking and of a similar illness 3 years previously. The illness responded promptly to treatment with amoxicillin, but her second cA2 infusion was withheld and the data for this patient are therefore not analyzed beyond week 2. Patient 18 showed significant bacteriuria on routine culture at week 6 (>10⁵/ml; lactose-fermenting coliform), but was asymptomatic. This condition also responded promptly to amoxicillin.

Routine analysis of blood samples showed no consistent adverse changes in hematologic parameters, renal function, liver function, or levels of C3, C4, or immunoglobulins during the 8 weeks of the trial. Four minor, isolated, and potentially adverse laboratory disturbances were recorded. Patient 2 experienced a transient rise in blood urea levels, from 5.7 mmol/liter to 9.2 mmol/liter (normal 2.5–7), with no change in serum creatinine. This change was associated with the temporary use of a diuretic, which had been prescribed for a non-rheumatologic disorder. The value normalized within 1 week and was classified as "probably not related" to cA2.

Patient 6 experienced a transient fall in the peripheral blood lymphocyte count, from 1.6×10^9 /liter to 0.8×10^9 /liter (normal 1.0–4.8). This abnormality was not seen at the next sample point (2 weeks later), was not associated with any clinical manifestations, and was classified as "possibly related" to cA2. Patients 10 and 18 developed elevated titers of anti-DNA antibodies at weeks 6 and 8 of the trial. Elevated anticardiolipin antibodies were also detected in patient 10. Both patients had a preexisting positive antinuclear antibody titer, and patient 10 had a history of borderline lymphocytopenia and high serum IgM. There were no clinical features of systemic lupus erythematosus, and the laboratory changes were judged "probably related" to cA2.

Efficacy of cA2. The pattern of response for each of the clinical assessments of disease activity and the derived IDA are shown in Table 2. All clinical assessments showed improvement following treatment with cA2, with maximal responses from week 3. Duration of morning stiffness decreased from a median of 180 minutes at study entry (week 0) to 5 minutes at week 6 ($P < 0.001$ by Mann-Whitney test, adjusted), representing a 97% improvement. The pain score decreased from 7.1 to 1.9 over the same period ($P < 0.001$, adjusted), representing an improvement of 73%. Similarly, the Ritchie Articular Index improved from 28 to 6 at week 6 ($P < 0.001$, adjusted; 79% improvement), and the swollen joint count decreased from 18




Figure 1. Swollen joint counts (maximum 28), as recorded by a single observer, in 20 patients with rheumatoid arthritis treated with cA2. The screening time point was within 4 weeks of entry to the study (week 0); data from patient 15 were not included after week 2 (dropout). Significance of the changes, relative to week 0, were determined by Mann-Whitney test (adjusted): $P > 0.05$ at week 1, $P < 0.02$ at week 2, $P < 0.002$ at weeks 3 and 4, and $P < 0.001$ at weeks 6 and 8. Bars show median values.

to 5 ($P < 0.001$, adjusted; 72% improvement). The individual swollen joint counts for all time points are shown in Figure 1.

Grip strength also improved; the median grip strength rose from 77 mm Hg (left) and 92 mm Hg (right) at week 0 to 119 (left) and 153 (right) at week 6 ($P < 0.04$ and $P < 0.05$, left and right hands, respectively; $P > 0.05$ both hands, adjusted for multiple comparisons). The IDA has a range of 1 (normal) to 4 (severe disease activity). The IDA showed a decrease from a median of 3 at study entry to 1.7 at week 6 ($P < 0.001$, adjusted). Patients were asked to grade their responses to cA2 using a 5-point scale. No patient recorded a response of "worse" or "no change" at any point in the trial. "Fair," "good," and "excellent" responses were classified as improvements of 1, 2, and 3 grades, respectively. At week 6, there was a median of 2 grades of improvement (Table 2).

We also measured changes in the patients' functional capacity, using the HAQ, as modified for British patients (range 0-3). The median (range) HAQ score improved from 2 (0.9-3) at study entry to 1.1 (0-2.6) by week 6 ($P < 0.001$ and $P < 0.002$ adjusted).

The changes in the laboratory values which reflect disease activity are shown in Table 3. The most rapid and impressive changes were seen in serum CRP levels, which fell from a median of 39.5 mg/liter at week 0 (normal < 10) to 8 mg/liter by week 6 of the trial ($P < 0.001$, adjusted), representing an improvement of

80%. Of the 19 patients with elevated CRP at study entry, 17 showed decreases to the normal range at some point during the trial. The improvement in CRP was maintained in most patients over the assessment period (Table 3 and Figure 2); the exceptions with high values at 4 and 6 weeks tended to be those with the highest starting values (data not shown).

The ESR also showed improvement, with a fall from 55 mm/hour at study entry (normal < 10 in males and < 15 in females) to 23 mm/hour at week 6 ($P < 0.03$ and $P > 0.05$ adjusted; 58% improvement). SAA levels were elevated in all patients at trial entry, and fell from a median of 245 mg/ml (normal < 10) to 58 mg/ml at week 1 ($P < 0.003$ adjusted; 76% improvement) and to 80 mg/ml at week 2 ($P < 0.04$, adjusted). No significant changes were seen in Hgb level, WBC count, or platelet count at week 6, although the platelet count did improve at weeks 2 and 3 compared with trial entry (Table 3).

The response data were also analyzed for each patient individually (not shown). The majority of patients had their best overall responses at week 6, at which time 13 assessed their responses as "good" while 6 assessed their responses as "fair." Eighteen of the 19 patients who completed the treatment schedule achieved an improvement in the IDA of 0.5 or greater at week 6, and 10 achieved an improvement of 1.0 or greater. All patients achieved a response at week 6




Figure 2. Serum C-reactive protein (CRP) levels (normal 0-10 mg/liter), as measured by nephelometry, in 20 patients with rheumatoid arthritis treated with cA2. The screening time point was within 4 weeks of entry to the study (week 0); data from patient 15 were not included after week 2 (dropout). Significance of the changes, relative to week 0, were determined by Mann-Whitney test (adjusted): $P < 0.001$ at week 1, $P < 0.003$ at week 2, $P < 0.002$ at week 3, $P < 0.02$ at week 4, and $P < 0.001$ at weeks 6 and 8. Bars show median values.

according to the index described by Paulus et al (35). At week 6, all patients showed a mean improvement of 30% or greater in the 6 selected measures of disease activity (see Patients and Methods), with 18 of the 19 patients showing a mean improvement of 50% or greater (data not shown).

Although the study was primarily designed to assess the short-term effects of cA2 treatment, followup clinical and laboratory data are available for those patients followed for sufficient time ($n = 12$). The duration of response in these patients, defined as the duration of a 30% (or greater) mean improvement in the 6 selected disease activity measures, was variable, ranging from 8 weeks to 25 weeks (median 14) (data not shown).

Comparison of the clinical and laboratory data for patients treated with 2 infusions of cA2 (each at 10 mg/kg) versus those treated with 4 infusions (each at 5 mg/kg) showed no significant differences in the rapidity or extent of response (data not shown).

Immunologic investigations and cytokines. Measurement of rheumatoid factor by RAPA showed 14 patients with significant titers ($\geq 1:160$) at trial entry. Of these, 6 patients showed a decrease of at least 2 titers on treatment with cA2, while the remaining patients showed a change of 1 titer or less. No patient showed a significant increase in rheumatoid factor titer during the trial (data not shown). The median titer in the 11 patients decreased from 1:2,560 at entry to 1:480 by week 8 ($P > 0.05$) (Table 3). Specific rheumatoid factor isotypes were measured by ELISA, and showed decreases in the 6 patients whose RAPA had declined significantly, as well as in some other patients (data not shown). Median values for the 3 isotypes in the 14 patients seropositive at trial entry were 119, 102, and 62 IU/ml (IgM, IgG, and IgA isotypes, respectively) and at week 8 were 81, 64, and 46 IU/ml ($P > 0.05$).

We tested sera from patients 1-9 for the presence of bioactive TNF, using the WEHI 164 clone 13 cytotoxicity assay (36). In 8 patients, serum samples spanning the entire trial period were tested, while for patient 9, only 3 samples (1 pretrial, 1 intermediate, and the last available sample) were tested. The levels of bioactive TNF were below the limit of sensitivity of the assay in the presence of human serum (1 pg/ml) (data not shown).

Since production of CRP and SAA are thought to be regulated in large part by IL-6, we also measured serum levels of this cytokine, using 2 different assays which measure total IL-6. In the Medgenix assay, IL-6 was significantly elevated in 17 of the 20 patients at

study entry. In this group, levels fell from 60 pg/ml (range 18-500) to 40 pg/ml (range 0-230) at week 1 ($P > 0.05$) and to 32 pg/ml (range 0-210) at week 2 ($P < 0.005$ and $P < 0.01$, adjusted). These results were supported by measurement of serum IL-6 in the first 16 patients in a separate ELISA developed in-house. IL-6 was detectable in 11 of these samples, with median (range) levels falling from 210 pg/ml (25-900) at entry to 32 pg/ml (0-1,700) at week 1 ($P < 0.02$ and $P < 0.04$, adjusted) and to 44 pg/ml (0-240) at week 2 ($P < 0.02$ and $P < 0.03$, adjusted).

We tested sera from patients 1-10 for the presence of antiglobulin responses to the infused chimeric antibody, but none were detected (data not shown). In many patients, however, cA2 was still detectable in serum samples taken at week 8 (data not shown) and this may have interfered with the ELISA.

DISCUSSION

This is the first report describing the administration of anti-TNF α antibodies for treatment of human autoimmune disease. Many cytokines are produced in rheumatoid synovium, but we chose to specifically target TNF α because of mounting evidence that it was a major molecular regulator in RA (21,22,26-28). The study results presented here support that view and allow 3 important conclusions to be drawn.

First, treatment with cA2 was safe and the infusion procedure was well tolerated. Although fever, headache, chills, and hemodynamic disturbance have all been reported following treatment with anti-CD4 or anti-CDw52 in RA (6,10), such features were absent in our patients. Also notable was the absence of any allergic event despite repeated treatment with the chimeric antibody, although the interval between initial and repeat infusions may have been too short to allow maximal expression of any antiglobulin response. The continuing presence of circulating cA2 at the conclusion of the trial may have precluded detection of antiglobulin responses, but also indicated that any such responses were likely to be of low titer and/or affinity. Although we recorded 2 episodes of infection among the study group, these were minor and the clinical courses were unremarkable. TNF α has been implicated in the control of *Listeria* and other infections in mice (37), but our limited experience does not suggest an increased risk of infection after TNF α blockade in humans.

The second conclusion concerns the clinical

efficacy of cA2. The patients we treated had longstanding, erosive, and for the most part, seropositive disease, and therapy with several standard DMARDs had failed. Despite this, the major clinical assessments of disease activity and outcome (morning stiffness, pain score, Ritchie articular index, swollen joint count, and HAQ score) showed statistically significant improvement, even after adjustment for multiple comparisons. All patients graded their response as at least "fair," with the majority grading it as "good." In addition, all achieved a response according to the criteria of Paulus et al and showed a mean improvement of at least 30% in 6 selected disease activity measures. The design of the trial does not allow these results to be attributed to the action of cA2 alone. However, the extent of the clinical improvements, their consistency throughout the study group, and the parallel changes in laboratory indices of disease activity (see below) are encouraging.

The improvements in clinical assessments following treatment with cA2 appear to be at least as good as those reported following treatment of similar patients with antileukocyte antibodies (6,10), although firm conclusions concerning each of these agents will require controlled, blinded studies. The two therapeutic approaches may already be distinguished, however, by their effects on the acute-phase response, since in several studies of antileukocyte antibodies, no consistent improvements in CRP or ESR were seen (4-6,8,10). In contrast, treatment with cA2 resulted in significant decreases in serum CRP and SAA values, with normalization of values in many patients. The changes were rapid and marked, and in the case of CRP, sustained for the duration of the study (Table 3). The decreases in ESR were less marked, achieving statistical significance only when unadjusted for the number of comparisons (Table 3).

These results are consistent with current concepts that implicate TNF α in the regulation of hepatic acute-phase protein synthesis, either directly, or by control of other, secondary, cytokines such as IL-6 (38,39). To investigate the mechanism of control of the acute-phase response in our patients, we measured serum TNF α and IL-6 before and after cA2 treatment. Bioactive TNF α was not detectable in sera obtained at baseline or subsequently. In view of previous reports of variability between different immunoassays in the measurement of cytokines in biologic fluids (40), we used 2 different assays for IL-6, and both demonstrated significant decreases in serum IL-6 levels by week 2. These findings support the other objective laboratory changes induced by cA2, and provide in

vivo evidence that TNF α may be a regulatory cytokine for IL-6 in this disease. Among the other laboratory tests performed, levels of rheumatoid factors fell significantly in 6 patients.

The mechanism of action of cA2 leading to the clinical responses outlined above was not established in this study. Neutralization of TNF α may have a number of beneficial consequences, including a reduction in the local release of cytokines such as IL-6 and other inflammatory mediators, and modulation of synovial endothelial/leukocyte interactions. cA2 may also bind directly to synovial inflammatory cells expressing membrane TNF α , with subsequent in situ cell lysis. Further studies should establish which actions of cA2 may be clinically important.

The results obtained in this small series have important implications, both scientifically and clinically. At the scientific level, the ability of the neutralizing antibody, cA2, to reduce acute-phase protein synthesis, reduce the production of other cytokines such as IL-6, and significantly improve the clinical state demonstrates that it is possible to interfere with the cytokine network in a useful manner without untoward effects. Due to the many functions and overlapping effects of cytokines such as IL-1 and TNF α , and the fact that cytokines induce the production of other cytokines and of themselves, there had been some pessimism as to whether targeting a single cytokine in vivo would have any beneficial effect (41,42). This view is clearly refuted. On the clinical side, the results of short-term treatment with cA2 are encouraging, and suggest that TNF α may be a useful new therapeutic target in RA.

ACKNOWLEDGMENTS

We thank Drs. R. McCloskey, M. Sanders, U. Nas-sander, C. Wortel, D. Zelinger, and R. Lorian of Centocor Inc. for their assistance with trial management. Dr. F. di Padova for the kind gift of anti-IL-6 monoclonal antibodies. Drs. J. Kirwan, H. Bird, and P. Schur for reviewing the manuscript, and the following physicians for the referral of patients included in this study: S. Allard, M. Corbett, C. Erhardt, A. Keat, C. Higgins, C. Mackworth-Young, G. Room, A. So, and P. Venables. We also thank Ms A. Hogg, Ms S. Tyler, and Ms G. Harris for excellent technical assistance and Dr. C. Plater-Zyberk for help in videotaping patient interviews.

REFERENCES

1. Pincus T, Callahan LF: Taking mortality in rheumatoid arthritis seriously: predictive markers, socioeconomic status and comorbidity. *J Rheumatol* 13:841-845, 1986

2. Scott DL, Symmons DPM, Coulton BL, Popert AJ: Long-term outcome of treating rheumatoid arthritis: results after 20 years. *Lancet* 1:1108-1111, 1987
3. Erhardt GC, Mumford PA, Venables PJW, Maini RN: Factors predicting a poor life prognosis in rheumatoid arthritis: an eight year prospective study. *Ann Rheum Dis* 48:7-13, 1989
4. Herzog C, Walker C, Muller W, Rieber P, Reiter C, Riethmüller G, Wassmer P, Stockinger H, Madic O, Pichler W: Anti-CD4 antibody treatment of patients with rheumatoid arthritis: I. Effect on clinical course and circulating T cells. *J Autoimmun* 2:627-642, 1989
5. Goldberg D, Morel P, Chatenoud L, Boitard C, Menkes CJ, Bertoye P, Revillard J, Bach J: Immunological effects of high dose administration of anti-CD4 antibody in rheumatoid arthritis patients. *J Autoimmun* 4:617-630, 1991
6. Horneff G, Burmester GR, Emmrich F, Kalden JR: Treatment of rheumatoid arthritis with an anti-CD4 monoclonal antibody. *Arthritis Rheum* 34:129-140, 1991
7. Kirkham BW, Pitzalis C, Kingsley GH, Chikanza IC, Sabharwal S, Barbatis C, Grahame R, Gibson T, Amlot PL, Panayi GS: Monoclonal antibody treatment in rheumatoid arthritis: the clinical and immunological effects of a CD7 monoclonal antibody. *Br J Rheumatol* 30:459-463, 1991
8. Reiter C, Kakavand B, Rieber EP, Schattenkirchner M, Riethmüller G, Krüger K: Treatment of rheumatoid arthritis with monoclonal CD4 antibody M-T151: clinical results and immunopharmacologic effects in an open study, including repeated administration. *Arthritis Rheum* 34:525-536, 1991
9. Wendling D, Wijdenes J, Racadot E, Morel-Fourrier B: Therapeutic use of monoclonal anti-CD4 antibody in Rheumatoid arthritis. *J Rheumatol* 18:325-327, 1991
10. Isaacs JD, Watts RA, Hazleman BL, Hale G, Keogen MT, Cobbold SP, Waldmann H: Humanised monoclonal antibody therapy for rheumatoid arthritis. *Lancet* 340:748-752, 1992
11. Ziff M: Rheumatoid arthritis: its present and future. *J Rheumatol* 17:127-133, 1990
12. Fontana A, Hengartner H, Weber E, Fehr K, Grob PJ, Cohen G: Interleukin-1 activity in the synovial fluid of patients with rheumatoid arthritis. *Rheumatol Int* 2:49-53, 1982
13. Buchan G, Barrett K, Turner M, Chantry D, Maini RN, Feldmann M: Interleukin-1 and tumour necrosis factor mRNA expression in rheumatoid arthritis: prolonged production of IL-1 α . *Clin Exp Immunol* 73:449-453, 1988
14. Chu CQ, Field M, Feldmann M, Maini RN: Localization of tumor necrosis factor α in synovial tissues and at the cartilage-pannus junction in patients with rheumatoid arthritis. *Arthritis Rheum* 34:1125-1132, 1991
15. Hirano T, Matsuda T, Turner M, Miyasaka N, Buchan G, Tang B, Sato K, Shimizu M, Maini R, Feldmann M, Kishimoto T: Excessive production of interleukin 6/B cell stimulatory factor-2 in rheumatoid arthritis. *Eur J Immunol* 18:1797-1801, 1988
16. Xu WD, Firestein GS, Taetle R, Kaushansky K, Zvaifler NJ: Cytokines in chronic inflammatory arthritis. II: Granulocyte-macrophage colony-stimulating factor in rheumatoid synovial effusions. *J Clin Invest* 83:876-882, 1989
17. Brennan FM, Zachariae COC, Chantry D, Larsen CG, Turner M, Maini RN, Matsushima K, Feldmann M: Detection of interleukin-8 biologic activity in synovial fluids from patients with rheumatoid arthritis and production of interleukin-8 mRNA by isolated synovial cells. *Eur J Rheumatol* 20:2141-2144, 1990
18. Brennan FM, Chantry D, Turner M, Foxwell B, Maini RN, Feldmann M: Detection of transforming growth factor- β in rheumatoid arthritis synovial tissue: lack of effect on spontaneous cytokine production in joint cell cultures. *Clin Exp Immunol* 81:278-285, 1990
19. Chu CQ, Field M, Abney E, Zheng RQH, Allard S, Feldmann M, Maini RN: Transforming growth factor- β 1 in rheumatoid synovial membrane and cartilage/pannus junction. *Clin Exp Immunol* 86:380-386, 1991
20. Kissonerghis AM, Maini RN, Feldmann M: High rate of HLA class II mRNA synthesis in rheumatoid arthritis joints and its persistence in culture: down regulation by recombinant interleukin-2. *Scand J Immunol* 29:73-82, 1989
21. Brennan FM, Chantry D, Jackson A, Maini RN, Feldmann M: Inhibitory effects of TNF α antibodies on synovial cell interleukin-1 production in rheumatoid arthritis. *Lancet* 2:244-247, 1989
22. Haworth C, Brennan FM, Chantry D, Turner M, Maini RN, Feldmann M: Expression of granulocyte-macrophage colony-stimulating factor (GM-CSF) in rheumatoid arthritis: regulation by tumour necrosis factor- α . *Eur J Immunol* 21:2575-2579, 1991
23. Brennan FM, Gibbons DL, Mitchell T, Cope AP, Maini RN, Feldmann M: Enhanced expression of tumour necrosis factor receptor mRNA and protein in mononuclear cells isolated from rheumatoid arthritis synovial joints. *Eur J Immunol* 22:1907-1912, 1992
24. Deleuran BW, Chu C-Q, Field M, Brennan FM, Mitchell T, Feldmann M, Maini RN: Localization of tumor necrosis factor receptors in the synovial tissue and cartilage-pannus junction in patients with rheumatoid arthritis: implications for local actions of tumor necrosis factor α . *Arthritis Rheum* 35:1170-1178, 1992
25. Keffer J, Probert L, Czirakis H, Georgopoulos S, Kaslaris E, Kioussis D, Kollias G: Transgenic mice expressing human tumour necrosis factor: a predictive genetic model of arthritis. *EMBO J* 10:4025-4031, 1991
26. Williams RO, Feldmann M, Maini RN: Anti-tumour necrosis factor ameliorates joint disease in murine collagen-induced arthritis. *Proc Natl Acad Sci U S A* 89:9784-9788, 1992
27. Feldmann M, Brennan FM, Chantry D, Haworth C, Turner M, Abney E, Buchan G, Barrett K, Barkley D, Chu A, Field M, Maini RN: Cytokine production in the rheumatoid joint: implications for treatment. *Ann Rheum Dis* 49:480-486, 1990
28. Brennan FM, Maini RN, Feldmann M: TNF α : a pivotal role in rheumatoid arthritis? *Br J Rheumatol* 31:293-298, 1992
29. Arnett FC, Edworthy SM, Bloch DA, McShane DJ, Fries JF, Cooper NS, Healey LA, Kaplan SR, Liang MH, Luthra HS, Medsger TA Jr, Mitchell DM, Neustadt DH, Pinals RS, Schaller JG, Sharp JT, Wilder RL, Hunder GG: The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 31:315-324, 1988
30. Mallia RK, Mace BEW: The assessment of disease activity in rheumatoid arthritis using a multivariate analysis. *Rheumatol Rehabil* 20:14-17, 1981
31. Steinbrocker O, Traeger CH, Batterman RC: Therapeutic criteria in rheumatoid arthritis. *JAMA* 140:659-662, 1949
32. Ritchie DM, Boyle JA, McInnes JM, Jasani MK, Dalakas TG, Grieverson P, Buchanan WW: Clinical studies with an articular index for the assessment of joint tenderness in patients with rheumatoid arthritis. *Q J Med* 147:393-406, 1968
33. Fuchs HA, Brooks RH, Callahan LF, Pincus T: A simplified twenty-eight-joint quantitative articular index in rheumatoid arthritis. *Arthritis Rheum* 32:531-537, 1989
34. Kirwan JR, Rebeck JS: Stanford health assessment questionnaire modified to assess disability in British patients with rheumatoid arthritis. *Br J Rheumatol* 25:206-209, 1986
35. Paulus HE, Egger MJ, Ward JR, Williams HJ, and the Cooperative Systematic Studies of Rheumatic Diseases Group: Analysis of improvement in individual rheumatoid arthritis patients treated with disease-modifying antirheumatic drugs, based on the findings in patients treated with placebo. *Arthritis Rheum* 33:477-484, 1990
36. Espevik T, Nissen-Meyer J: A highly sensitive cell line, WEHI 164 clone 13, for measuring cytotoxic factor/tumour necrosis

- factor from human monocytes. *J Immunol Methods* 95:99-105, 1986
37. Havell EA: Evidence that tumour necrosis factor has an important role in antibacterial resistance. *J Immunol* 143:2894-2899, 1989
38. Fong Y, Tracey KJ, Moldawer LL, Hesse DG, Manogue KB, Kenney JS, Lee AT, Kuo GC, Allison AC, Lowry SF, Cerami A: Antibodies to cachectin/tumour necrosis factor reduce interleukin-1 β and interleukin-6 appearance during lethal bacteremia. *J Exp Med* 170:1627-1633, 1989
39. Guerne P-A, Zuraw BL, Vaughan JH, Carson DA, Lotz M: Synovium as a source of interleukin-6 in vivo: contribution to local and systemic manifestations of arthritis. *J Clin Invest* 83:585-592, 1989
40. Roux-Lombard P, Steiner G, and the Cytokine Consensus Study Group Of The European Workshop For Rheumatology Research: Preliminary report on cytokine determination in human synovial fluids: a consensus study of the European Workshop for Rheumatology Research. *Clin Exp Rheumatol* 10:515-520, 1992
41. Kingsley G, Panayi G, Lanchbury J: Immunotherapy of rheumatic diseases: practice and prospects. *Immunol Today* 12:177-179, 1991
42. Trentham DE: Immunotherapy and other novel therapies. *Curr Opin Rheumatol* 3:369-372, 1991

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.